


GlucoBalance®

Support for Normal Glucose Levels



Effective regulation of blood glucose has important implications for health. Even mild disruptions of glucose homeostasis can have adverse consequences. Chronic diabetes may result in cardiovascular disease, neuropathy, blindness, or renal failure. Hypoglycemia (also called reactive hypoglycemia or dysinsulinism), though not generally associated with the organ damage seen in diabetes, can be responsible for a number of troublesome physical and psychological symptoms.

The human body possesses a complex set of checks and balances to maintain blood glucose concentrations within a narrow range. Blood sugar control is influenced by the pituitary, thyroid, and adrenal glands, as well as by the pancreas, liver, kidney, and even skeletal muscle.

Glucose homeostasis also depends on the presence of a wide range of micronutrients. In the typical American diet, high in refined and processed foods, many of these micronutrients are in short supply. In addition, some individuals with blood sugar disorders may have a special dietary need for higher amounts of one or more micronutrients.

The following nutrients are particularly important when considering blood sugar disorders:

Formulated for Biotics Research by Jonathan V. Wright, M.D. and Alan R. Gaby, M.D., leaders in the field of nutritional science.

Wide clinical use in the U.S.A. 1,000 mcg chromium. Chromium is essential for glucose metabolism.

To place your order for **GlucoBalance®** or for additional information please contact us:

(800) 231-5777

Utilizing "The Best of Science and Nature" to Create Superior Nutritional Supplements

6801 Biotics Research Drive • Rosenberg, TX 77471
biotics@bioticsresearch.com • www.bioticsresearch.com

These statements have not been evaluated by the Food and Drug Administration. These products are not intended to diagnose, treat, cure, or prevent any disease.

 **BIOTICS**
RESEARCH
CORPORATION
*Utilizing "The Best of Science and Nature"
to Create Superior Nutritional Supplements*

LIT-011 © Copyright 2007-16 Rev. 11/16

Chromium—The effect of chromium on glucose metabolism apparently requires its conversion to glucose tolerance factor (GTF), a low-molecular-weight compound that contains chromium, niacin (nicotinic acid), glycine, glutamic acid, and cysteine. GTF, has been shown to potentiate the action of insulin at the cellular level.^(1,4)

Tissue chromium levels were found to decline with age in Americans.⁽²⁾ In other studies, including one by the U.S. Department of Agriculture, more than 50% of people consumed less than the lower level of chromium recommended by the National Academy of Sciences, Nutritional Research Council.^(3,37) Chromium aspartate is a well-utilized form of supplemental

Contains both Niacin, a vital component of GTF (Glucose Tolerance Factor) & Niacinamide, necessary for producing NAD (Nicotinamide Adenine Dinucleotide) in insulin secreting pancreatic B cells.

Formulated with a uniquely high amount of biotin- 3,000 mcg. Biotin is essential for glucose phosphorylation by hepatic glucokinase, the first step in glucose utilization.

Antioxidant activity in vitamin E and selenium.

chromium being solubilized at a wide range of pH. The amounts of chromium used in most clinical trials (*150 to 200 ug/day) are apparently inadequate for some patients, even when more efficient Chromium compounds are used. Larger amounts of chromium, such as 500 to 1,000 ug/day, have often had a greater benefit.⁽⁴⁾

Niacin and

Niacinamide—As a component of glucose tolerance factor, niacin plays an important role in carbohydrate metabolism. Many refined foods consumed by Americans are depleted of niacin. Grains and other foods that are “enriched” usually contain added niacinamide, which apparently cannot be converted by the human body into niacin. In addition, many vitamin supplements contain niacinamide, rather than niacin. Although niacinamide is capable of performing most of the functions of vitamin B₃, a small amount of niacin seems to be necessary for the synthesis of GTF.⁽⁵⁾ Both niacin and niacinamide may also be important for blood sugar control through a mechanism unrelated to GTF. As precursors to NAD, which is an important metabolite concerned with intracellular energy production, niacin and niacinamide may prevent the depletion of NAD in pancreatic B cells.

Biotin—The initial step in glucose utilization by the cell is its phosphorylation, mediated by the biotin-dependent enzyme hepatic glucokinase. Thus adequate biotin intake is required to initiate intracellular glucose into the cell.⁽⁶⁾

Biotin may also play a role in stabilizing blood sugar levels through biotin-dependent enzymes acetyl CoA carboxylase and pyruvate carboxylase.⁽³⁷⁾ Thus biotin deficiency should be avoided in those with blood sugar disorders.

Pyridoxine (vitamin B₆)—Serum vitamin B₆ levels were below normal in 25% of 518 diabetics.⁽⁷⁾ Particularly where peripheral neuropathy is present, the inadequate B₆ intake should be contemplated.⁽⁸⁾

Copper—Because the typical American diet contains only about half of the RDA (2mg/day) for copper^(9,36) deficiency of this mineral may be common. Copper is involved with insulin binding, and copper deficiency in mammals may be reflected in increased glycosylated hemoglobin, indicative of chronically raised blood sugar levels.⁽¹⁰⁾

Magnesium—The American diet is often low in magnesium. Dietary surveys have shown that 80-85% of American women consume less than the RDA for the mineral.⁽¹¹⁾ Daily magnesium intake in two other studies was only about two thirds of the RDA.^(12,13,36) This may be particularly relevant in diabetics, where magnesium deficiency is thought to play a role in the development of insulin resistance.⁽¹⁴⁾ Serum magnesium has been found to be significantly lower in many diabetics,⁽¹⁵⁾ therefore, it is reasonable to make sure that diabetics have adequate dietary intake of magnesium. Low magnesium levels may also be associated with hypoglycemia.⁽¹⁶⁾

Zinc—Zinc is involved both with insulin synthesis by pancreatic B cells⁽¹⁷⁾ and insulin binding to liver and adipose tissue cells.^(18,19) People with zinc deficiency may have significantly higher glucose levels and lower insulin levels than similar patients without zinc deficiency.⁽²⁰⁾

Vitamin C (ascorbic acid)—Ascorbic acid levels may be lower in diabetics than controls⁽²²⁾ and patients with inadequate ascorbic acid levels may be found to have abnormal blood sugar curves.⁽²³⁾ In addition, ascorbic acid may compete with glucose for transport across cell membranes,⁽²⁴⁾ so that in hyperglycemia, ascorbic acid transport into the cell may be inhibited.⁽²⁵⁾

Ascorbic acid deficiency may allow sorbitol to accumulate in erythrocytes, which may pre-dispose diabetics to certain types of end-organ damage.⁽²⁶⁾ Care should be taken to ensure that those with blood sugar abnormalities have adequate ascorbic acid intake.

Manganese — Manganese is a cofactor for certain enzymes involved in the intermediary metabolism of carbohydrates. In addition, the concentration of manganese in the pancreas is approximately ten times higher than in other organs.⁽²⁷⁾

The optimal intake of manganese is not known but at least half of the manganese is lost when whole grains are replaced by refined flour.⁽²⁹⁾ The American diet may be low in manganese.⁽³⁶⁾

Both B₁₂ and folate are involved in a number of different steps in carbohydrate metabolism and the incidence of B₁₂ deficiency was significantly greater in a series of diabetics than in the general population.⁽³⁰⁾ Folate is involved with gluconeogenesis as a cofactor with key enzymes in the liver and small intestine.⁽³¹⁾ Deficiency of either B₁₂ or folate is to be avoided in those with blood sugar abnormalities.

Vitamin B1 (thiamin) — Central to carbohydrate metabolism and Krebs Cycle function is adequate thiamin levels, and diabetics are more often deficient in thiamin.⁽³²⁾ Care should be taken to ensure that thiamin intake is adequate in those with diabetes and hypoglycemia.

Carnitine — Carnitine is involved as part of a vital transport mechanism of fat metabolism in which fat enters energy production pathway. Carnitine supplementation may be considered in those with blood sugar abnormalities.

Vanadium — Vanadate is an oxidized form of vanadium. Due to possible insulinotropic effects of vanadate⁽³³⁾, inadequate amounts of this trace mineral is undesirable in those with blood sugar abnormalities.

Vitamin E and Selenium — Vitamin E and selenium are essential nutritional factors which act as antioxidants and may be involved in glucose balance.^(34,35) As many complications associated with diabetes may be related to excess free radical activity, prudence demands that adequate selenium and vitamin E be supplied in the diabetic diet.

References:

1. Toepfer EW, Mertz W, Polansky MM, Roginski EE, Wolf WR. Preparation of chromium- containing material of glucose tolerance factor activity from brewer's yeast extracts and by synthesis. *J Agric Food Chem* 1977;25:162-166.
2. Schroeder HA, Nason AP, Tipton IH. Chromium deficiency as a factor in atherosclerosis. *J Chronic Dis* 1970;23:123-142.
3. Anderson RA, Kozlovsky AS. Chromium intake, absorption and excretion of subjects consuming self-selected diets. *Am J Clin Nutri* 1985;41:1177-1183.
4. Glinsmann WH, Mertz W. Effect of trivalent chromium on glucose tolerance. *Metabolism* 1966;15:510-502.
5. Urberg, M, Zemel MB. Evidence for synergism between chromium and nicotinic acid in the control of glucose tolerance in elderly humans. *Metabolism* 1987;36:896-899.
6. Anonymous. Biotin and glucokinase in the diabetic rat. *Nutr. Rev* 1970;28:242-244.
7. Davis RE, Calder JS, Curnow DH. Serum pyridoxal and folate concentrations in diabetics. *Pathology* 1976;8:151-156.
8. Jone CL, Gonzalez V. Pyridoxine deficiency: a new factor in diabetic neuropathy. *J. Am Podiatry Assoc* 1978;68:646-653.
9. Wolf WR, Holden J, Greene FE. Daily intake of zinc and copper from self selected diets. *Fed Proc* 1977;36:1175.
10. Klevay LM, Canfield WK, Gallagher SK, Henrickson LK, Bolonchuk W, et al. Diminished glucose tolerance in two men due to a diet low in ic acid in the control of glucose tolerance in elderly humans. *Metabolism* 1987;36:896-899.
11. Morgan KJ, Stampely GL, Zabik ME, Fischer DR. Magnesium and calcium dietary intakes of the U.S. population. *J Am Coll Nutr* 1985;4:195-206.
12. Lakshmanan FL, Rao RB, Kim WW, Kelsay JL. Magnesium intakes, balances and blood levels of adults consuming self-selected diets. *Am J Clin Nutri* 1984;40:1380-1389.
13. Srivastava US, Nadeau MH, Gueneau L. Mineral intakes of university students; magnesium content. *Nutr Rep Int* 1978;18:235-242.
14. Yajnik CS, Smith RF, Hockaday TDR, Ward NI. Fasting plasma magnesium concentrations and glucose disposal in diabetes. *Br Med J* 1984;288:1027-1028.
15. Ceriello A, Guigliano D, Dello Russo P, Passariello N. Hypomagnesemia in relation to diabetic retinopathy. *Diabetes Care* 1982;5:558-559.
16. Stebbing JB, Turner MO, Franz KB. Reactive hypoglycemia and magnesium. *Magnesium Bull* 1982;2:131=134.



BIOTICS
RESEARCH
CORPORATION



Utilizing "The Best of Science and Nature"
to Create Superior Nutritional Supplements

GlucoBalance®

Support for Normal Glucose Levels

17. Emdin SO, Dodson GG, Cutfield JM, Cutfield SM. Role of zinc in insulin biosynthesis. Some possible zinc-insulin interactions in the pancreatic B-cell. *Diabetologia* 1980;19:172-182.
18. Arquilla ER, Packer S, Tarmas W, Miyamoto S. The effect of zinc on insulin metabolism. *Endocrinology* 1978;103:1440-1449.
19. Herington AC. Effect of zinc on insulin binding to rat adipocytes and hepatic membranes and to human placental membranes and IM-9 lymphocytes. *Horm Metab Res* 1985;17:328-332.
20. Wolman SL, Anderson GH, Marliss EB, Jeejeebhoy KN. Zinc in total parental nutrition: requirements and metabolic effects. *Gastroenterology* 1979;76:458-467.
21. Holden JM, Wolf WR, Mertz W. Zinc and copper in self-selected diets. *J Am Diet Assoc* 1979;75:23-28.
22. Sarji KE, Kleinfelder J, Brewington P, Gonzalez J, Hempling H, et al. Decreased platelet vitamin C in diabetes mellitus: possible role in hyperaggregation. *Thromb Res* 1979;15:636-650.
23. Secher K. The bearing of the ascorbic acid content of the blood on the course of the blood sugar curve. *Acta Med Scand* 1942;60:255-265.
24. Verlangieri AJ, Sestito J. Effect of insulin on ascorbic acid uptake by heart endothelial cells: possible relationship to retinal atherogenesis. *Life Sci* 1981;29:5-9.
25. Hutchinson ML, Lee WYL, Chen MS, Davis KA, Ely JTA, et al. Effects of glucose and select pharmacologic agents on leukocyte ascorbic acid levels. *Fed Proc* 1983;42:930.
26. Vinson JA, Staretz ME, Bose P, Kassim HM, Basalyga BS. *In vitro* and *in vivo* reduction of erythrocyte sorbitol by ascorbic acid. *Diabetes* 1989;38:1036-1041.
27. Fore H. Manganese-induced hypoglycemia. *Lancet* 1963;1:274-275.
28. Shrader RE, Everson GJ. Pancreatic pathology in manganese-deficient guinea pigs. *J Nutr* 1968;94:269.
29. Wenlock RW, Buss DH, Dixon EJ. Trace nutrients. Manganese in British food. *Br J Nutr* 1979;41:253-261.
30. Wilkinson JF. Diabetes mellitus and pernicious anemia. *Br Med J* 1963;1:676-677.
31. Lifshitz F, Moak SA, Wapnir Ra. Folic acid in the prevention of fasting induced hypoglycemia. *Pediatr Res* 1977;11:518.
32. Haugen HN. The blood concentration of thiamin in diabetes. *Scand J Clin Lab Invest* 1964;16:260-266.
33. Fagin JA, Ikejiri K, Levin SR. Insulinotropic effects of vanadate. *Diabetes* 1987;36:1448.
34. Asayama K, Kooy NW, Burr IM. Effect of vitamin E deficiency and selenium deficiency on insulin secretory reserve and free radical scavenging systems in islets: decrease of islet manganosuperoxide dismutase. *J Lab Clin Med* 1986; 107:459-464.
35. Echert CD, Breskin MW, Wise WW, Knopp RH. Association between low serum selenium and diminished visual function in diabetic women. *Fed Proc* 1985;44:1670.
36. Pennington JAT, Young BE, Wilson DB, Johnson RD, Vanderveer JE. Mineral content of foods in total diet: selected minerals in food survey, 1982-1984. *Jour Am Diet Assoc*. 86:876, 1986.
37. Coggeshall JC, Heggars JP, Robson MC, Baker H. Biotin status and plasma glucose in diabetics. *Ann NY Acad Sci* 1985;447:389-393.

GlucoBalance® is available in 90-count (#1850) and 180-count bottles (#1851).

Supplement Facts

Serving Size: 6 Capsules
Servings Per Container: 15

	Amount Per Serving	% Daily Value		Amount Per Serving	% Daily Value
Vitamin A (as retinyl acetate)	1,500 mcg RAE	167%	Zinc (as zinc picolinate and zinc citrate)	30 mg	272%
Vitamin C (as calcium ascorbate and ascorbic acid)	500 mg	556%	Selenium (as selenomethionine)	150 mcg	272%
Vitamin D (as cholecalciferol)	2.5 mcg	13%	Copper (as copper gluconate)	2 mg	222%
Vitamin E (as d-alpha tocopheryl acetate)	270 mg	1,800%	Manganese (as manganese citrate)	20 mg	870%
Thiamin (B1) (as thiamin mononitrate)	50 mg	4,167%	Chromium (as chromium aspartate)	1,000 mcg	2,857%
Riboflavin (B2)	25 mg	1,923%	Potassium (as potassium aspartate)	99 mg	2%
Niacin (as niacinamide and niacin)	150 mg	938%	Vanadium (as vanadium aspartate)	20 mcg	*
Vitamin B6 (as pyridoxine hydrochloride)	30 mg	1,765%	L-Carnitine fumarate	30 mg	*
Folate (as calcium folinate)	800 mcg DFE	200%	* Daily Value not established		
Vitamin B12 (as methylcobalamin)	50 mcg	2,083%	Other ingredients: Capsule shell (gelatin and water) and magnesium stearate (vegetable source).		
Biotin	3,000 mcg	10,000%	This product is gluten, dairy and GMO free.		
Pantothenic Acid (as calcium pantothenate)	100 mg	2,000%	RECOMMENDATION: Six (6) capsules per day, two (2) capsules taken with each meal as a dietary supplement or as otherwise directed by a healthcare professional.		
Calcium (as ascorbate, citrate and carbonate)	200 mg	15%	WARNING: If you are taking Diabetes medications, do not use this product without medical supervision.		
Magnesium (as aspartate, citrate and oxide)	400 mg	95%			

KEEP OUT OF REACH OF CHILDREN

Store in a cool, dry area.
Sealed with an imprinted safety seal for your protection.

Product # 1850 Rev. 11/16

(800) 231-5777

6801 Biotics Research Drive • Rosenberg, TX 77471
biotics@bioticsresearch.com • www.bioticsresearch.com



BIOTICS
RESEARCH
CORPORATION



Utilizing "The Best of Science and Nature" to Create Superior Nutritional Supplements